

Low-Level Laser Therapy (780 nm) on VEGF Modulation at Partially Injured Achilles Tendon

Julio Fernandes de Jesus, MSc,¹ Diva Denelle Spadacci-Morena, PhD,¹ Nayra Deise dos Anjos Rabelo, MSc,² Carlos Eduardo Pinfildi, PhD,³ Thiago Yukio Fukuda, PhD,⁴ and Helio Plapler, PhD⁵

Abstract

Objective: The aim of this study was to analyze the modulatory effects of near infrared (780 nm) low-level laser therapy (LLLT) on the presence of the vascular endothelial growth factor (VEGF) in the partially injured Achilles tendons of rats. **Background:** LLLT stimulates the healing process for Achilles tendon injuries, although the extent of the modulatory effect of LLLT on the VEGF levels found in the injured tendons remains unclear. **Methods:** Sixty-five male Wistar rats were distributed in the following seven groups: LASER 1, 3, and 7 (10 partially injured Achilles tendons in each group, which were treated with LLLT for 1, 3, and 7 days, respectively); Sham 1, 3, and 7 (same injury, with simulated LLLT); Control group containing the five remaining animals and in which no procedures were performed. LLLT was applied once a day for 10 sec, with a mean power of 70 mW and fluency of 17.5 J/cm². After euthanasia, all of the Achilles tendons were surgically removed and the VEGF levels were analyzed using immunohistochemistry. **Results:** The VEGF levels remained close to normal ($p > 0.05$) when comparing the experimental groups (LASER and Sham: 1, 3, and 7) with the Control group. **Conclusion:** LLLT did not stimulate the expression of VEGF in the treated Achilles tendons.

Introduction

DESPITE THE FACT that Achilles tendon is the largest and strongest tendon in the human body, it exhibits high rates of injuries (acute and degenerative), which can take weeks or even months to heal. This slow healing process has been correlated with morphological abnormalities, proinflammatory factors triggered by the injuries, and the vascular characteristics of the Achilles tendon.^{1–6}

The vascular endothelial growth factor (VEGF) is one of the most important factors involved in angiogenesis, exhibiting a notable presence in Achilles tendon injuries.⁷ This increase in the VEGF has been interpreted as beneficial, since it is an attempt to increase the number of blood vessels and the nutritional contribution. Consequently, the proliferative rate increases, as does the healing process in the Achilles tendon.^{8,9}

However, there is evidence that this strategy is ineffective, as it is not strong enough to heal the Achilles tendon and the new vessels that appear are accompanied by new nerve bundles, thereby increasing nociceptive afferents and pain.¹⁰ In addition, VEGF induces the expression and

presence of distinct deleterious and degrading factors in the tendon tissue, including matrix metalloproteinases (MMP), which delay the healing process and may even weaken the injured tendon further.^{8,11}

Low-level laser therapy (LLLT) has provided satisfactory results when used to treat Achilles tendon injuries, due to the proliferative and modulatory stimulation it promotes.^{3–6,12} There are also some reports of increases in the expression and presence of VEGF in several experimental situations in which LLLT was involved. This phenomenon—if not clarified and controlled—could hinder the healing process of the Achilles tendon.^{8,13–16}

Although there are positive reports of the interaction between LLLT and Achilles tendon injuries, there are still uncertainties about the main active pathways involved and whether this angiogenic characteristic of LLLT, which is found in different clinical and experimental situations, also occurs in Achilles tendon injuries.^{3,5,6} Thus, the aim of the present study was to analyze the modulatory effects of near infrared (780 nm) LLLT on the presence of VEGF in the partially injured Achilles tendons of rats.

¹Laboratory of Physiopathology, Instituto Butantan, São Paulo, Brazil.

²Rehabilitation Sciences Program, Universidade Nove de Julho-UNINOVE, São Paulo, Brazil.

³Plastic Surgery Department, Universidade Federal de São Paulo-UNIFESP, São Paulo, Brazil.

⁴Knee and Hip Rehabilitation, Instituto Trata, São Paulo, Brazil.

⁵Surgery Department, Universidade Federal de São Paulo-UNIFESP, São Paulo, Brazil.

Materials and Methods

The present study was approved by the Research Ethics Committee of the *Universidade Federal de São Paulo* under protocol number 0074/2011.

Sixty-five male 12-week-old Wistar rats (*Rattus norvegicus*: var. *albinus*, *Rodentia*, *Mammalia*), weighing between 270 and 300 g, were used. They were kept in standard polypropylene cages, with a light/dark cycle of 12 h, a temperature of $\sim 20^{\circ}\text{C}$, and 65% of humidity. They received water and rat chow *ad libitum*.

Injury procedure

Sixty animals were randomly anesthetized with an intraperitoneal injection of ketamine hydrochloride (100 mg/kg) and xylazine hydrochloride (50 mg/kg). The five remaining animals were not submitted to anesthesia.

After anesthesia, manual trichotomy of the right hind paw was performed (Achilles tendon areas). The animals were positioned at an injury device base—a mini-guillotine dispositive designed to drop weight and promote injury by direct trauma. The right hind paw was positioned on the equipment base, and mild traction was exerted through ankle dorsiflexion until the dorsal region of the paw rested against the injury device base. Finally, a weight of 186 g was released from a height of 20 cm onto the Achilles tendon of each animal, corresponding to a potential energy of 364.9 mJ at the time of trauma.^{3,4}

Group distribution

After partially injuring the Achilles tendon of 60 animals, they were distributed randomly into six groups:

- Sham 1: 10 tendons submitted to simulated LLLT for 1 day.
- LASER 1: 10 tendons submitted to LLLT application for 1 day.
- Sham 3: 10 tendons submitted to simulated LLLT for 3 days.
- LASER 3: 10 tendons submitted to LLLT application for 3 days.
- Sham 7: 10 tendons submitted to simulated LLLT for 7 days.
- LASER 7: 10 tendons submitted to LLLT application for 7 days.

The five remaining animals (10 tendons) were placed at the Control group and were not submitted to any procedures.

Laser treatment

The AsGaAl LLLT (MM Optics[®], Model: Twin Laser) equipment was used. Table 1 presents the LLLT parameters applied.

Laser was performed manually once a day at the central portion of injured Achilles tendons, with the animal gently caged and immobilized during treatment and/or simulation period. The animals were treated for 1 day (LASER 1), 3 consecutive days (LASER 3), or 7 consecutive days (LASER 7). Treatment started 1 min after induced injury and contact technique and continuous emission were used in all irradiations.^{3,4,17}

TABLE 1. LLLT PARAMETERS

| <i>LLLT parameters</i> | |
|--|-----------------------------|
| <i>Light source</i> | <i>Laser diode (AsGaAl)</i> |
| Wavelength, nm | 780 |
| Mean power (Pm), mW | 70 |
| Beam area, cm ² | 0.04 |
| Power density (ΔP), W/cm ² | 1.75 |
| Energy density (ΔE), J/cm ² | 17.5 |
| Duration of each treatment session, sec | 10 |
| Dose of each treatment session, J | 0.7 |
| Cumulative dose (LASER 1, 3, and 7, respectively), J | 0.7, -2.1, -4.9 |

LLLT, low-level laser therapy.

In Sham groups (1, 3, and 7), simulated applications of LLLT were performed, with the apparatus turned off. However, contact was made between the equipment and the injured area for the same period of time as in the treated groups.

Euthanasia

The animals in the LASER 1/Sham 1, LASER 3/Sham 3, and LASER 7/Sham 7 groups were euthanized (inhalational anesthetic overdose with isoflurane, 5% for induction and 2% maintenance until euthanized) 24 h, 4 days, and 8 days after the injury had been induced, respectively. The animals in the Control group were euthanized on the last day of the experiment (eighth day).

Sample processing

After euthanasia, tendons of the two hind paws were surgically removed from animals in the Control group, as well as the 60 tendons (right hind paw) of animals belonging to the other groups.

The tendons were immediately washed in saline solution (0.9%) and then fixed in paraformaldehyde (4%) in a Millonig 0.1 M buffer (pH 7.2–7.4) for 12 h.

After fixation, the tendons were washed in the buffer and dehydrated in increasing solutions of ethyl alcohol (from 70% to 100%) for 45 min in each solution. Next, they were diaphanized in xylene (two baths of 45 min each) and included in paraffin, positioned in such a way that longitudinal slices could be obtained. Microtome slices (model: RM2155; Leica) with a thickness of 5 μm were used in immunohistochemistry. The slides were silanized (3-aminopropyl-triethoxysilane—Sigma) for slice adhesion.

Each slide was prepared with a slice from each tendon to quantify the VEGF levels, a process that was repeated thrice. Therefore, three nonconsecutive slides were obtained for each animal studied.

All of the sections were dewaxed, hydrated, and submitted to the above method.

Analysis of angiogenesis (immunohistochemistry)

A specific antibody kit for the VEGF was used (anti-VEGF monoclonal; brand: Abcam—AB1316; dilution:

1/200). The dewaxed slices were washed in distilled water and Tris-phosphate buffer pH 7.4 (TBS).

Antigen retrieval was performed using pork trypsin 0.25% in phosphate buffer (code T7409; Sigma Aldrich) for 20 min at 37°C. Endogenous peroxidase was blocked with hydrogen peroxide 10v (3%). The samples were then washed in tap water, deionized water, and then left in phosphate buffer pH 7.4 (PBS). Nonspecific proteins were blocked through the immersion of the slides in casein (Synth) and diluted in PBS (pH 7.4) for 5 min at room temperature.

The slides were incubated with the primary antibodies in a specific diluent (Spring Bioscience) for 24 h at 4°C and then incubated with the secondary antibody Histofine (Nishirei Biosciences, Inc.) for 45 min at 37°C. Subsequently, the slides were washed in water and the sections were placed in diaminobenzidine (DAB) (Spring Biosciences) for 10 min at room temperature.

Finally, the slices were counterstained with Harris hematoxylin (Merck) and mounted with Entellan (Merck).

The effect of the treatment on the VEGF response of the injured Achilles tendon was analyzed in three different anatomical regions of the tendon (proximal, intermediary, and distal) to identify and quantify positive cell types. Two random images of each of the anatomical regions analyzed were collected, with a zoom of 200× (objective 20×), using a DFC420 camera (Leica Microsystems) coupled to a DMLB microscope (Leica Microsystems). After obtaining the images, cells that exhibited positive markings were identified and quantified using version 4.5.0 of the LAS computerized image system (Leica Microsystems).

Statistical analysis

The Statistical Package for Social Sciences (SPSS®—version 15.0) was used, with the level of significance set at 5% ($p < 0.05$). The Levene's tests and one-way analysis of variance (ANOVA) were used to assess the homogeneity of the data and to compare the results between the groups, respectively. The data were displayed in mean and standard deviation values and the result of the comparisons was reported using the F test, when homogeneity of the variances was recorded. If differences were to be recorded between the groups, Tukey's multiple comparisons tests (*post hoc*) were conducted.

Results

Table 2 presents the total number of positive cells for VEGF in the proximal, intermediary, and distal regions analyzed. The experimental groups (LASER and Sham: 1, 3, and 7) did not exhibit elevated levels of angiogenesis at the analyzed regions and also at the total average when compared with the Control group ($p > 0.05$).

The intergroup analysis (Sham 1, 3, and 7 vs. LASER 1, 3, and 7) concerning the number of applications (1, 3, and 7) and the regions analyzed (proximal, intermediary, and distal) also exhibited no statistically significant differences for the levels of VEGF ($p > 0.05$), with similar angiogenesis patterns (Fig. 1).

Discussion

The aim of this controlled laboratory study was to analyze the effect of LLLT on the VEGF modulation of the

TABLE 2. TOTAL NUMBER OF CELLS (MEAN \pm SD) POSITIVE FOR THE VEGF

| | Proximal | Intermediary | Distal | Mean |
|---------|-----------------|-----------------|-----------------|-----------------|
| Control | 16.2 \pm 3.4 | 17.6 \pm 6.1 | 18.8 \pm 6.7 | 17.5 \pm 2.4 |
| Sham 1 | 22.5 \pm 8.4 | 24.4 \pm 11.7 | 18.4 \pm 8.9 | 21.8 \pm 7.7 |
| LASER 1 | 21.6 \pm 8.6 | 26.4 \pm 8.2 | 23.4 \pm 12.0 | 23.8 \pm 8.1 |
| Sham 3 | 25.7 \pm 10.8 | 24.3 \pm 10.4 | 24.6 \pm 15.0 | 24.9 \pm 9.4 |
| LASER 3 | 28.1 \pm 21.0 | 24.3 \pm 14.3 | 24.5 \pm 11.0 | 25.6 \pm 13.0 |
| Sham 7 | 23.0 \pm 10.4 | 25.1 \pm 12.4 | 16.7 \pm 6.1 | 21.6 \pm 7.7 |
| LASER 7 | 23.3 \pm 8.7 | 24.6 \pm 12.4 | 23.4 \pm 5.9 | 23.8 \pm 4.7 |

VEGF, vascular endothelial growth factor.

partially injured Achilles tendons of rats. The most significant result obtained was the maintenance of the quantity of VEGF at near-normal levels in the tendons treated with LLLT.

Achilles tendon injuries are complex and multifactorial, as is the subsequent healing process.^{4,18,19} LLLT has provided promising results when used in the treatment of Achilles tendon injuries, basically acting in two pathways: proliferative (regeneration and collagen reorganization) and modulatory (controlling the inflammatory process).³⁻⁶

Besides the regeneration and reorganization of collagen fibers and the control of the inflammatory process, modulating the VEGF levels is important when seeking to repair injured Achilles tendons, given that its high and uncontrolled presence may hamper and delay the healing process once VEGF induces the expression of tendon tissue degrading factors and also may increase pain (stimulating the presence of new nerve bundles) during rehabilitation programs.^{8,10,11,18,20}

There are reports in the literature of increases in VEGF levels based on the induction of Achilles tendon injuries. However, unlike the present one, these studies used different methods of tendon injury (collagenase), which in turn may be more specific to trigger the modulation of VEGF levels.²⁰ In addition, the periods studied were longer than those proposed in this study, which may have perpetuated the tendon injury for enough time to induce and favor the detection of VEGF modulations.²⁰

The LLLT applied in this study, different several previous reports in the literature,^{13-16,21} did not cause an increase in VEGF levels, which may be a positive effect given the complications that this increase can cause when found in injured tendon tissues.⁷⁻¹¹ Although the active mechanisms of LLLT in Achilles tendon injuries have not been completely clarified, it is probable that this LLLT (which exhibits anti-inflammatory effects) reduces the migration of proinflammatory agents (macrophages and monocytes) to the injury site, thereby maintaining near-normal VEGF levels, since VEGF is also secreted by these agents.^{6,11,19,20}

Furthermore, the interaction between LLLT and the different types of tissues and factors involved in the healing/regeneration processes depends on a number of factors: wavelength, mean power, and the dose and frequency of the treatment.²²⁻²⁴ There are reports in the literature showing modulations in VEGF levels with a lower mean power (22 mW) and a higher final energy (1.54 J)²⁰ than those used in the present study. It is probable that our dosimetric




FIG. 1. Photomicrography of the mean values for cells that were positive (✓) for the VEGF (immunohistochemistry) at 20×: (A) Control, (B) Sham 1, (C) LASER 1, (D) Sham 3, (E) LASER 3, (F) Sham 7, and (G) LASER 7. VEGF, vascular endothelial growth factor.

parameters did not achieve the required specificity and thus did not promote an increase in the VEGF levels.

Conversely, there are also reports in the literature showing the proliferation and reorganization of collagen fibers, as well as the reduction of the levels of proinflammatory and deleterious factors of injured Achilles tendons using the same LLLT protocol (wavelength, dose, and treatment frequency). These reports also used the same experimental model as the present study, demonstrating the specificity of this LLLT in the interaction with these other factors during the repair of injured Achilles tendons.^{3–6,18}

Therefore, this LLLT protocol exhibited positive effects on the healing of injured Achilles tendons (specifically by increasing collagen proliferation and organization) and on the control of the inflammatory process, acting in a proliferative

and modulatory, rather than an angiogenic pathway, since there was no increase in the VEGF levels after its application.

Although no VEGF modulation was found with the LLLT used in this study, uncertainty remains about its effects when applied in other experimental models (more aggressive and chronic injuries) and other treatment protocols—since this research investigated a small period and a single method of analysis—and there is also uncertainty about the best LLLT parameters.

Conclusions

Near infrared 780 nm LLLT did not stimulate VEGF expression at the partially injured Achilles tendons at the early stages of the repair process.

Acknowledgment

The authors thank the São Paulo Research Foundation (FAPESP) for supporting this study (2011/00979-1) and Magna Aparecida Maltauro Soares for providing technical support.

Author Disclosure Statement

No competing financial interests exist.

References

- Doral MN, Alam M, Bozkurt M, et al. Functional anatomy of the Achilles tendon. *Knee Surg Sports Traumatol Arthrosc* 2010;18:638–643.
- O'Brien M. The anatomy of the Achilles tendon. *Foot Ankle Clin* 2005;10:225–238.
- Wood VT, Pinfield CE, Neves MA, Parizoto NA, Hochman B, Ferreira LM. Collagen changes and realignment induced by low-level laser therapy and low-intensity ultrasound in the calcaneal tendon. *Lasers Surg Med* 2010;42:559–565.
- Oliveira FS, Pinfield CE, Parizoto NA, et al. Effect of low level laser therapy (830 nm) with different therapy regimes on the process of tissue repair in partial lesion calcaneus tendon. *Lasers Surg Med* 2009;41:271–276.
- de Jesus JF, Spadacci-Morena DD, Rabelo ND, Pinfield CE, Fukuda TY, Plapler H. Low-level laser therapy on tissue repair of partially injured achilles tendon in rats. *Photomed Laser Surg* 2014;32:345–350.
- de Jesus JF, Spadacci-Morena DD, dos Anjos Rabelo ND, Pinfield CE, Fukuda TY, Plapler H. Low-level laser therapy in IL-1 β , COX-2, and PGE2 modulation in partially injured Achilles tendon. *Lasers Med Sci* 2015;30:153–158.
- Petersen W, Pufe T, Kurz B, Mentlein R, Tillmann B. Angiogenesis in fetal tendon development: spatial and temporal expression of the angiogenic peptide vascular endothelial cell growth factor. *Anat Embryol (Berl)* 2002; 205:263–270.
- Pufe T, Petersen WJ, Mentlein R, Tillmann BN. The role of vasculature and angiogenesis for the pathogenesis of degenerative tendons disease. *Scand J Med Sci Sports* 2005; 15:211–222.
- Pufe T, Petersen W, Kurz B, Tsokos M, Tillmann B, Mentlein R. Mechanical factors influence the expression of endostatin—an inhibitor of angiogenesis—in tendons. *J Orthop Res* 2003;21:610–616.
- Alfredson H, Ohberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 2003;11:334–338.
- Wang H, Keiser JA. Vascular endothelial growth factor upregulates the expression of matrix metalloproteinases in vascular smooth muscle cells: role of flt-1. *Circ Res* 1998; 83:832–840.
- Fukuda TY, Tanji MM, Jesus JF, Sato MN, Duarte AJ, Plapler H. Single session to infrared low level diode laser on TNF- α and IL-6 cytokines release by mononuclear spleen cells in mice: a pilot study. *Lasers Surg Med* 2010;42:584–588.
- Szymanska J, Goralczyk K, Klawe JJ, et al. Phototherapy with low-level laser influences the proliferation of endothelial cells and vascular endothelial growth factor and transforming growth factor-beta secretion. *J Physiol Pharmacol* 2013;64:387–391.
- Martignago CC, Oliveira RF, Pires-Oliveira DA, et al. Effect of low-level laser therapy on the gene expression of collagen and vascular endothelial growth factor in a culture of fibroblast cells in mice. *Lasers Med Sci* 2014;30:203–208.
- de Oliveira TS, Serra AJ, Manchini MT, et al. Effects of low level laser therapy on attachment, proliferation, and gene expression of VEGF and VEGF receptor 2 of adipocyte-derived mesenchymal stem cells cultivated under nutritional deficiency. *Lasers Med Sci* 2015;30:217–223.
- Cury V, Moretti AI, Assis L, et al. Low level laser therapy increases angiogenesis in a model of ischemic skin flap in rats mediated by VEGF, HIF-1 α and MMP-2. *J Photochem Photobiol B* 2013;125:164–170.
- Koshima H, Kondo S, Mishima S, et al. Expression of interleukin-1 β , cyclooxygenase-2, and prostaglandin E2 in a rotator cuff tear in rabbits. *J Orthop Res* 2007;25:92–97.
- Marcos RL, Arnold G, Magnenet V, Rahouadj R, Magdalou J, Lopes-Martins RÁ. Biomechanical and biochemical protective effect of low-level laser therapy for achilles tendinitis. *J Mech Behav Biomed Mater* 2014;29:272–285.
- Pires D, Xavier M, Araújo T, Silva JA Jr, Aimbire F, Albertini R. Low-level laser therapy (LLLT; 780 nm) acts differently on mRNA expression of anti- and pro-inflammatory mediators in an experimental model of collagenase-induced tendinitis in rat. *Lasers Med Sci* 2011;26:85–94.
- Casalechi HL, Leal-Junior EC, Xavier M, et al. Low-level laser therapy in experimental model of collagenase-induced tendinitis in rats: effects in acute and chronic inflammatory phases. *Lasers Med Sci* 2013;28:989–995.
- Park IS, Mondal A, Chung PS, Ahn JC. Prevention of skin flap necrosis by use of adipose-derived stromal cells with light-emitting diode phototherapy. *Cytherapy* 2015;17: 283–292.
- Bjorndal J, Coupe C, Ljunggren A. Low level laser therapy for tendinopathy. Evidence of dose-response pattern. *Phys Ther Rev* 2001;6:91–99.
- Karu T. Photobiology of low-power laser effects. *Health Phys* 1989;56:691–704.
- Carrinho PM, Renno AC, Koeke P, Salate AC, Parizotto NA, Vidal BC. Comparative study using 685-nm and 830-nm lasers in the tissue repair of tenotomized tendons in the mouse. *Photomed Laser Surg* 2006;24:754–758.

Address correspondence to:
 Diva Denelle Spadacci-Morena
 Laboratory of Physiopathology
 Instituto Butantan
 São Paulo
 SP 05504-900
 Brazil

E-mail: divamorena@butantan.gov.br